

Received: 12 January 2014 • Accepted: 01 February 2014

Short C

doi:10.15412/J.JBTW.01030103

Study on traditional training methods based on Health Belief Model on exercise performance in patients with Myocardial Infarction

Ladan EbrahimPouriyan¹, Mohammad Mahboubi^{2*}, Zeynab Karimi³, Mahmoud Kazemi⁴

¹ MSc in Medical Surgical Nursing, Kermanshah University of Medical Sciences, Kermanshah, Iran

² PhD in Health Services Administration, Kermanshah University of Medical Sciences, Kermanshah, Iran

³ MS.c Student in Department of Health Education & Health promotion Tehran University of Medical Sciences, Tehran, Iran

⁴ MS.c Student in Medical Education, Kermanshah University of Medical Sciences, Kermanshah, Iran

*correspondence should be addressed to Mohammad Mahboubi, PhD in Health Services Administration, Kermanshah University of Medical Sciences, Kermanshah, Iran; Tell: +989181323095; Fax: +98; Email: mm59m@yahoo.com.

ABSTRACT

In Iran, Deaths due to coronary artery diseases is very increasing. And unfortunately every year due to lack of physical activity and decreased cardio activity the age group of this disease is decreasing in a way that now this disease is seen in the youth, And one of basic tools is changing the patients lifestyle in the educational program as a part of the treatment plan. This study has been done with the aim to compare the traditional training methods based on Health Belief Models on exercise performance in patients with myocardial infarction. In this *Quasi- Experimental* Research 74 patients with myocardial infarction were chosen by purposefully sampling method then they were chosen randomly in both training and control groups, Training Group achieved education based on the concepts and components of health beliefs model training aimed at increasing the Perceived severity of patients and Control group achieved traditional education. The health belief model of education based on the performance of activity in the experimental group than the control group found no significant difference ($P = 0.000$). The perceived benefits of physical activity after intervention using the Student T test with a mean of 46.19 in the test group and the average 94.17 in the control group was significant. Education on the Health Belief Model in patients with myocardial infarction increases the perceived severity of patients and improvement in performance activity.

Key words: Health Belief Model, performance activity, myocardial infarction

Copyright © 2014 Ladan EbrahimPouriyan et al. This is an open access article distributed under the Creative Commons Attribution License.

1. INTRODUCTION

Myocardial infarction is the most common health problem threatening human life in the world. About one and a half million cases of myocardial infarction occur in the United States of America, and every 30 seconds one is diagnosed with myocardial infarction (1, 2). Every 10 minutes, one person loses his life due to myocardial infarction, and more than 50% of patients will need rehabilitation measures following myocardial infarction (3). It is estimated that by 2030, 32.5% of deaths would be

caused by cardiovascular disease in Europe. Cardiovascular disease has also become a social problem in Iran. The disease accounted for the largest share with 4 deaths of every 6 deaths per 10,000 of population. Unfortunately, the age of onset of cardiovascular diseases has declined in Iran (4) and hence now, they are seen not only in the middle ages, but also in the young people. Accurate estimates of the prevalence of risk factors of cardiovascular diseases are unattainable; however, the prevalence of identified risk factors has changed by increased awareness among the patients and diet changes as well as

changes in the patient's lifestyle (5, 6). It was observed in many studies conducted in order to detect risk factors on the incidence of cardiovascular disease in women(2010), That the risk of cardiovascular diseases is less in women with more physical activity who sit only for 4 hours a day than in women with low physical activity, It was mentioned in countless review articles that inactivity has a considerable role among risk factors for cardiovascular diseases with adverse effects on the progression of coronary atherosclerosis in men and women, and found that exercise may reduce the risk of coronary diseases (7). Considering the overwhelming evidence, the presence of relationship between the individuals' lifestyle and developing of cardiovascular diseases, the necessity of emphasis on individuals' lifestyle as an important factor in prognosis and prevention of recurrent cardiovascular diseases have been demonstrated. It should be noted that regular activity in patients who experienced heart attack lead to reduced risk of cardiac arrhythmias and improved perfusion through the coronary arteries and increased blood sub-flow to the heart and enables the person to return to the status before illness (8). Also, physical inactivity is the causing factor in 15% of chronic diseases, including heart disease (7). One of the most needed care services for cardiac patients after myocardial infarction is providing a developed educational program regarding compliance (9) in daily exercise program and lifestyles. In many countries, there are programs to encourage the community toward a Healthy Heart through education and counseling and encouraging the public and the patients to reduce risk factors such as sedentary lifestyle. Different approaches are required to change the behavior and beliefs of the society individuals, one of (10) them is the effective model on health education and behavior change of the Health Belief Model, the "Health Belief Model" was first introduced in 1950 by a number of psychologists. Based on this model, an individual adopts a preventive behavior when he affected by following factors, including:

- ✓ Perceived susceptibility: Understanding and believing in being exposed to the risk of disease
- ✓ Perceived severity: Understanding and believing in the seriousness of the problem that the problem can

lead to death or other serious consequences for the individual

- ✓ Perceived barriers: Physical, psychological or financial barriers against adopting hygiene practices
- ✓ Perceived benefits: Person's belief achieved in analysis of behavior advantages
- ✓ Guiding instructions: Including acceptable health messages, mass communication and individuals capable of influencing on taking measures of the target group
- ✓ Taking measures: Observing targeted health behavior influenced by above structures.

Health advises to adopt healthy behaviors, are only practiced by the patient and his family if they have become as belief in the patient's thought, mind and practice. For implementation of these recommendations by the patient and his family their participation in the care and preservation of healthy behaviors are must. Their participation can be realized only if such guidelines have become as their health and hygiene beliefs (11). Given the sufficient studies in connection with evaluation of the knowledge about coronary arteries diseases, proper exercise activities, perceived severity of illness, perceived benefits and barriers resulting from proper exercising in patients with myocardial infarction have not been conducted, This study was performed aiming at comparison the effects of two methods of training by traditional method and method based on Health Belief Model on functional activities of patients with myocardial infarction.

2. MATERIALS AND METHODS

This study was performed by a quantitative and experimental method. The study population included patients with

myocardial infarction hospitalized in Imam Ali (AS) and Imam Reza (AS) health centers affiliated to Kermanshah University of Medical Sciences in 2013. The study samples included 74 patients with inclusion criteria (Literacy level as reading and writing, aged between 18 and 70 years, no underlying disease with 48 hours passed from MI). The purposive-based sampling was done randomly in two experimental and (12) controlled groups. The sample size was estimated using Moradi's research methodology model for 74 cases. Data collection tool included a 61-item questionnaire based on health belief model and a researcher-made checklist. The questionnaire used in this study was developed according to the questionnaire designed in Moradi's research regarding the application of health belief model in adoption of healthy behaviors in patients undergoing coronary artery surgery (12). Based on research subject and in accordance with experts' opinions and the research goals, some changes were made. To acquire dace and content validity of the questionnaire and the checklist, the opinions and considerations of nine faculty members of Kermanshah Nursing and Midwifery School and Two cardiologists were collected, and their corrective feedbacks were applied. The questionnaire consisted of 12 demographic questions, 13 questions on the knowledge about the disease and how to exercise properly, 11 questions on perceived severity of the disease and 20 questions related to perceived benefits and barriers resulting from doing exercise activities and 5 questions with regard to the practice manuals. The checklist included 21 questions: 6 questions on demographic characteristics of family members of the patient and 15 questions about the

patient's performance in regard to the exercise activities. The Cronbach's coefficient alpha method was used to measure the reliability of the questionnaire and the checklist. Thus, 12 participants completed the questionnaire and the checklist in interviews within a 15-day interval. Finally, the reliability was obtained as $r = 0.81$. Data was analyzed using descriptive and inferential statistical tests related to the study by using SPSS software, version 18.

3. RESULTS AND DISCUSSION

The findings showed that there was no statistically significant difference between the samples of two test and control groups in variables of age, sex, employment status, previous history of stroke and death un the family and the first-degree relatives and knowledge score of each dimension ($p < 0.05$). In both test and control groups, the highest percentage frequency was related to men with 73% of and 65.3%, respectively. The highest frequency percentages in the experimental group and control group were related to the high school degree and guidance school, respectively, that using the K-2 test, no significant difference was seen between the two groups concerning education level ($P = 0.266$). In terms of employment status, the highest percentage frequency in both test and control groups were related with employees for 24.3% and 23.3% in values. The paired t-test results for the areas of knowledge aspects are given in Table 2 .

Table 1 . Average rating of knowledge studied groups before and after intervention

Dimensions of Consciousness		Control	Case	P- value
		M±SD	M±SD	
Awareness about CHD	Before	96.1±23.1	2±21.1	0.789
	After	61.4±67.1	90.2±24.1	0.000
Awareness of physical Activity	Before	75.6±98.1±	86.6±77.1	0.236
	After	99.8±47.1	67.6±27.2	0.000
The Total Score	Before	83.10±77.2	82.10±27.3	0.907

After	41.18±24.4	75.1±24.2	0.000
-------	------------	-----------	-------

The findings also showed that the trainings based on the health belief model in the test group has caused statistically significant increase in awareness rate of the disease ($p = 0.002$), perceived severity ($p = 0.003$), perceived benefits resulting from adherence to diet and exercise ($p = 0.001$), perceived barriers due to doing diet and exercise ($p = 0.000$) as well as improved exercise performance of the samples ($p = 0.004$). Although the training increased the patient's nutritional

performance, however, this increase was not significant statistically (Table 2).

Table 2 . Comparative study of concepts and components of the health belief model studied groups before and after intervention

Dimension		Control	Case	P- value
		M±SD	M±SD	
Awareness	Before	43.71±67.9	86.86±46.14	0.907
	After	91.4±18.13	44.13±11.18	00.000
Perceived Severity	Before	78.25±55.9	79.25±37.10	0.862
	After	49.36±9.7	60.27±41.10	0.000
Perceived Benefits	Before	77.43±38.3	70.44±15.7	0.734
	After	61.47±36.4	49.43±44.3	0.000
Perceived Barriers	Before	37.85±89.4	86.01±52.18	0.804
	After	53.1±35.16	85.16±5.01	0.000
Function	Before	31.8±57.8	46.09±76.12	0.047
	After	50.04±91.1	45.01±48.17	0.000

The results showed that training in the framework of health belief model leads to increased knowledge in understanding rate of heart stroke disease and increased perceived benefits in patients with heart failure, improves the activities functionality of the patients as the impact of such education. The study results confirmed the results of Zigheymat studies concerning the application of the health belief model in patients undergoing coronary surgery and the results of Ali and Haddad studies on application of this model in sports participation rates

in patients with MI in Jordan (2, 13). However, in a study conducted by Abedi et al "the effect of lifestyle changes using health belief model on cardiac risk factors in postmenopausal women in Ahwaz", it was found that all the components of Health Belief Model, except for the knowledge about cardiovascular diseases, were improved in the intervention group compared to the control group. The researcher stated that to increase awareness, training based on age and education level of the patients is required. The results

of this research study were not consistent with the results of the conducted study regarding increased knowledge of the patients about the disease (14). Sanaei et al conducted a study entitled as "Effects of family-oriented training on the level of adherence to the exercise program in patients undergoing coronary artery bypass surgery. It was found that after educational intervention based on health belief model in the experimental group, the quality of their adherence to the exercise program improved. One of the most important barriers in achieving compliance with treatment programs is because of the absence of the patient's relatives. One can also say that one of the most important factors in being influenced by the surgery guidelines model is the presence or absence of the relatives that the results of the present study are consistent with Sanaei et al. study. Also, the results of this study were consistent with Gucci study, in which by use of health belief model, the role of exercise in American women was evaluated. In his study, after the intervention, a significant difference in the exercise performance of two test and control groups was observed (14). The research findings showed a significant relationship between age and physical performance; however, there was no significant difference between gender and level of knowledge in the two groups. This finding was not consistent with the study of Motamedi et alon examining the impact of education based on the health belief model on the prevention of cutaneous leishmaniasis. The reason for this difference is caused by the age range difference of the studied patients in the two studies (11). No significant differences were found between the samples gender and the patients performance in both test and control groups, while the study of women's knowledge and performance regarding nutrition and exercise and its relation with cardiovascular diseases indicated the failure of women to realize the trainings (15, 16).

4. CONCLUSION

Given the positive results and impact of using Health Belief Model in educating patients with myocardial infarction and the impact of the model on patients' lifestyle changes, it is recommended to use the model in rehabilitation programs in patients with MI. The Community vulnerability to heart diseases and enormous cost of treating the diseases with lack of consistent and systematic plan for training the people at risk, lack of emphasis on its prevention prior to treatment policy is practiced by the Ministry of Health and other involved

organizations, lack of full belief of the society members about the role of nutrition and exercise pattern in the prevention of diseases due to local justifications and promoting of some superstitious miraculous medicines through illegal networks advertising are only a part of the problems faced by the health society in Iran. Lack of practical plans in this regard will bring a serious threat in the next decade to the community health.

ACKNOWLEDGMENT

No mentioned by authors.

AUTHORS CONTRIBUTION

This work was carried out in collaboration between all authors.

CONFLICT OF INTEREST

Authors have declared that no conflict interests exist.

REFERENCES

1. Rescuers A, Cpr LRA, Bls HP, Therapies E, Resuscitation N. Highlights of the 2010 American Heart Association Guidelines for CPR and ECC.
2. Ebrahimi PL, Salehi S, Pourmirza KR, Abdeyazdan G, Sharifi A. ASSESSING THE EFFECT OF HEALTHY BELIEF MODEL APPLICATION ON BEHAVIOR CHANGE OF THE PATIENTS WITH MYOCARDIAL INFARCTION.
3. Shidfar MR, Hosseini M, Zadeh DS, Asasi N, Majlesi F, Nazemi S. Effectiveness of an educational program on knowledge and attitudes of angina patients in Mashhad, Iran: results of an intervention. *Journal of Birjand University of Medical Sciences*. 2007;14(1):9-15.
4. Vahedian Azimi A, Alhani F, Ahmadi F, Kazemnejad A. Effect of family-centered empowerment model on the life style of myocardial infarction patients. *Iranian Journal of Critical Care Nursing*. 2009;2(4):127-32.
5. Kitkungvan D, Lynn Fillipon NM, Dani SS, Downey BC. Low-density lipoprotein cholesterol target achievement in patients at high risk for coronary heart disease. *Journal of clinical lipidology*. 4(4):293-7.
6. Tan YY, Gast Gc Fau - van der Schouw YT, van der Schouw YT. Gender differences in risk factors for coronary heart disease. (1873-4111 (Electronic)).
7. reprint from UpToDate O, Aroesty JM, Gersh BJ, Saperia GM. FRCP, MACC.
8. Manson JE, Greenland P Fau - LaCroix AZ, LaCroix Az Fau - Stefanick ML, Stefanick MI Fau - Mouton CP, Mouton Cp Fau - Oberman A, Oberman A Fau - Perri MG, et al. Walking compared with vigorous exercise for the prevention of cardiovascular events in women. (1533-4406 (Electronic)).
9. Corrigan PW. Enhancing personal empowerment of people with psychiatric disabilities. *American Rehabilitation*. 2004;28(1):10-21.
10. Ebrahim S, Taylor F, Ward K, Beswick A, Burke M, Davey Smith G. Multiple risk factor interventions for primary prevention of coronary heart disease. *Cochrane Database Syst Rev*. 1(1).
11. Motamedi N, Hejazi SH, Hazavehei SMM, Saberi S, Rahimi E. Effect of education based on Health Belief Model on promoting preventive behavior of coetaneous leishmaniasis. *MilMed Journal*. 11(4):231-6.
12. Moradi N. Application of health belief model in examining Tasyrbkar adopt health-enhancing behaviors in patients undergoing coronary artery bypass graft in the center of Imam Ali (AS) in Kermanshah: MS Thesis]. University Baghiatolla, 2006.[Persian].
13. Ali AL, Hadad LG. The effect of the health belief model in explaining exercise participation among Jordanian myocardial infarction patient. *Journal Transculate Nurse*. 15(2):114-210.
14. Huang MSL, Kandiah M, Yassin Z, Abedi P. Lifestyle change using the health belief model to improve cardiovascular risk factors among

postmenopausal women. Scientific Research Journal of Health System Research.7(1):127-37.

15.Koch J. The role of exercise in the African-American woman with type 2 diabetes mellitus: application of the health belief model. (1041-2972 (Print)).

16.Avazeh A, Jafari N, Rabie Siahkali S, Mazloomzadeh S. Knowledge level attitude and performance of women on diet and exercise and their relation with cardiovascular diseases risk factors. JOURNAL OF

ZANJAN UNIVERSITY OF MEDICAL SCIENCES AND HEALTH SERVICES.